

Alfred-Nobel-Str. 10 • 40789 Monheim • Germany Phone: +49 (0)2173 993730 contact@cube-biotech.com www.cube-biotech.com

PureCube NHS-Activated Agarose XL

| Product | Catalog No. | Package size |
|--|-------------|-----------------------|
| PureCube NHS-Activated Agarose XL (10 mL) | 58303 | 20 mL 50% suspension |
| PureCube NHS-Activated Agarose XL (50 mL) | 58305 | 100 mL 50% suspension |
| PureCube NHS-Activated Agarose XL (250 mL) | 58310 | 500 mL 50% suspension |

Product Description

PureCube NHS activated Agarose XL has been synthesized for the direct covalent coupling of biomolecules via free amine groups.

This activated chromatography matrix is based on custom-made agarose particles, consisting of 6% cross-linked agarose. The material is highly porous to allow for optimal protein interaction. Cross-linked agarose is also physically very stable, making it suitable for purification processes under low pressure with flow rates of up to 6 mL/min (optimal 0.5–2 mL/min). Our agarose is very homogeneous in size with a medium particle diameter of 400 μm , yielding a high degree of reproducibility between individual purification runs.

A NHS (N-hydroxy succinimide)-activated carboxy function is coupled to the magnetic agarose with a C6 spacer to obtain a matrix with highest binding capacity for amino functions. The carboxylic group density is higher than 15 μ mol/ml, as determined by acidimetric titration. The binding capacity of NHS-activated Agarose was determined with GFP protein to be up to 20 mg/ml.

PureCube NHS activated Agarose XL is delivered as a 50% suspension. Therefore, 2 mL suspension will yield a 1 mL bed volume. The suspension contains anhydrous isopropanol to prevent microbial growth and reduce hydrolysis.

Note: It is highly recommended to withdraw resin under protective gas and after equilibration to room temperature in order to avoid condensation of humidity and hydrolysis. Furthermore, it is recommended to directly react the resin with the target molecule after withdrawal, because hydrolysis will reduce activated groups directly after contact with humidity.

Shipping & Storage

| Shipment Temperature | Ambient temperature |
|----------------------|--|
| Storage | 100% Isopropanol, anhydrous, or another aprotic anhydrous solvent. |

Additional Information

For coupling protocols, and protocols for protein purification, please visit our webpage at: www.cube-biotech.com/protocols. For affinity purification of tagged proteins, Cube Biotech offers dedicated agarose resins, magnetic beads and prepacked cartridges. Also available are a range of ultrapure detergents and buffers for extraction and purification of proteins. See www.cube-biotech.com/products for details.

<u>Disclaimer</u>: Our products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

Proteins are our passion.