

Phone: +49 (0)2173 993730 contact@cube-biotech.com www.cube-biotech.com

PureCube Compact Cartridge Zn-NTA 1 mL & 5 mL

Product	Catalog No.	Package size
PureCube Compact Cartridge Zn-NTA (1 x 1 mL)	31602-Zn	1 x 1 mL prepacked column
PureCube Compact Cartridge Zn-NTA (5 x 1 mL)	31604-Zn	5 x 1 mL prepacked columns
PureCube Compact Cartridge Zn-NTA (1 x 5 mL)	31606-Zn	1 x 5 mL prepacked column
PureCube Compact Cartridge Zn-NTA (5 x 5 mL)	31608-Zn	5 x 5 mL prepacked columns

Product Description

The PureCube Compact Cartridge **Zn**-NTA is a chromatography column prepacked with PureCube Zn-NTA Agarose. The column is stored in buffer containing 20% ethanol to prevent microbial growth. The PureCube Compact Cartridge Zn-NTA is available in two sizes, 1 mL bed volume and 5 mL bed volume (dimensions given in Product Specifications). Both column sizes exhibit excellent chemical resistance to most commonly used reagents and the End Plugs include standard connections compatible with common chromatography instruments (such as ÄKTA). The 5 mL column has two layers of mesh (coarse and fine) at one end to give excellent flow distribution. The void volume in each End Plug is minimal, because the fluid is introduced through a narrow flow path (i.e. 1 mm hole).

Product Specifications

Parameter	PureCube Compact Cartridge Zn-NTA, 1 mL	PureCube Compact Cartridge Zn-NTA, 5 mL
Functional Group	Zn-NTA	z n-NTA
Dimensions [mm]	8.0 x 35	17 x 35
Column Body Material	Polypropylene	Polypropylene
End Plug Material	Polypropylene	Polypropylene
Inlet/Outlet	10-32 UNF female thread	10-32 UNF female thread
Matrix	7.5% highly cross-linked agarose	7.5% highly cross-linked agarose
Particle Diameter	40 μm	40 μm
Protein Binding Capacity*	Up to 70 mg	Up to 350 mg
Max. Flow Rate	6 mL/min	6 mL/min
Recommended Flow Rate**	0.5-2.0 mL/min	0.5-2.0 mL/min
Recommended Operational Pressure	Up to 5 bar (72 psi)	Up to 3 bar (42 psi)
pH Stability	2-14	2-14

^{*} Protein binding capacity can vary for different proteins

Affinity Resin

PureCube Zn-NTA Agarose was developed for the affinity purification of proteins carrying a polyhistidine tag. This affinity chromatography matrix is based on BioWorks Workbeads, consisting of 7.5% crosslinked agarose. The material is highly porous to allow for optimal protein interaction. Cross-linked agarose is also physically very stable, making it suitable for purification processes under low pressure with variable flow rates. Our agarose is very homogeneous in size with a medium particle diameter of 40 µm, yielding a high degree of reproducibility between individual purification runs.

An NTA ligand is coupled to the agarose resin and carefully loaded with zinc ions to obtain a matrix with highest binding capacity for histidine residues. The metal ion capacity is about 15 μ eqv Zn²⁺/mL. Other possible metal ions are Co²⁺, Ni²⁺, Fe³⁺, Al³⁺, resulting in different affinities, e.g., for his-tagged proteins or phosphorylated proteins. If required, the zinc ions can be removed from the agarose matrix using five wash steps with 100 mM EDTA, and the matrix can be recharged with a different metal ion. Alternatively, please contact us for unloaded NTA agarose. NTA matrices charged with other metals are available upon request.

Dynamic Protein Binding Capacity

t should be considered that the dynamic binding capacity strongly correlates with flow rate and other parameters such as protein size and buffer conditions. It is recommended to use the lowest flow rate possible to achieve highest binding capacity.

Compatibility

For cleaning purposes, PureCube Zn-NTA Agarose is very stable and can resist the following conditions in most situations: All commonly used aqueous buffers, pH 2-14, 100% methanol, 100% ethanol, 8 M urea, 6 M guanidinium hydrochloride, 30% (v/v) acetonitrile, and up to 10 mM DTT.

Shippping and Storage

Parameter	PureCube Compact Cartridge Zn-NTA, 1 mL	PureCube Compact Cartridge Zn-NTA, 5 mL
Shipment Temperature	Ambient temperature	Ambient temperature
Storage Buffer	20% ethanol, pH 6.5	20% ethanol, pH 6.5
Storage Temperature	2-8 °C	2-8 °C

Additional Information

For protein purification and cleaning protocols, including protocols for packing chromatography columns, please visit our webpage at www.cube-biotech.com/protocols. For purification of his-tagged proteins from dilute solutions, we recommend using PureCube Zn-NTA MagBeads. For affinity purification of GST-tagged, rho1D4-tagged or strep®-tagged proteins, Cube Biotech offers dedicated agarose resins, magnetic beads and pre-packed cartridges. Also available are a range of ultrapure detergents and buffers for the extraction and purification of proteins. See www.cube-biotech.com/products for details.

<u>Disclaimer</u>: Our products are intended for molecular biology applications. These products are not intended for the diagnosis, prevention, or treatment of a disease.

Trademarks: Strep-tag® (IBA GmbH).